

After Surgery Care Instructions

Thank you for having your pet spayed or neutered by the APF Community Spay/Neuter clinic. The APF is an independent nonprofit organization that receives no government funding. It is only through community support and donations that we are able to provide this service.

Your pet had major surgery with general anesthesia and will need to be cared for afterwards to help avoid complications. Following the instructions below will help ensure your pet has a successful recovery. Animals may be groggy or sleepy from the anesthesia. Your pet may be a little agitated or aggressive due to the after-effects of anesthesia. Avoid handling the animal too much as he/she may try to bite or scratch you. Your pet should be isolated from children and other pets. Your pet may also have poor balance; this will make getting out of the car and climbing stairs more difficult.

- **Limit Activity** We strongly recommend you keep your pet confined in a crate or small room the night after surgery. Keep your pet confined to a warm, quiet environment and separate from other pets and young children for the first 12 hours. No running, jumping, playing, rough housing, bathing or swimming or strenuous activity for 7 to 10 days following surgery. When outdoors, dogs should be kept on a short leash.
- **Feeding** Anesthesia tends to make animals experience nausea, so your pet may not want to eat when he/she gets home after surgery. Re-introduce food slowly. For adult animals offer ½ the normal amount of food and water when your pet is fully awake. If vomiting occurs, wait until the next day to give more food. Do not change your pet's diet at this time and do not offer table foods. Your pet's appetite should return gradually within 24 hours of surgery.
- **Under 4 months of age** Animals under 4 months of age should eat right away and may receive the normal amount of food and water.
- **Incision Site** What you see on the day of surgery is what we consider normal. The incision site should be checked twice daily. Check for excessive redness, swelling, discharge, blood or opening of the incision site. A very small amount of redness/swelling at incision may occur. Do not clean or apply any topical ointment to the incision site as this may cause the surgical glue to dissolve too quickly. You must keep your pet from licking or biting at the incision site as this could cause the wound to reopen and become infected. If your pet is excessively licking we recommend you purchase an E-collar or one of the topical ointments that you can spray near the incision site. (Place a clean finger over the incision site and spray around the area. **DO NOT SPRAY DIRECTLY ONTO THE INCISION - IT WILL STING**). Any breakdown (opening) of the incision, discharge, or drainage should be reported.
- **Female dogs and cats** will have a mid-line incision to their abdomen where the uterus and ovaries were removed. This incision is sealed with internal sutures that will dissolve. Surgical glue may have also been applied to the skin to seal the incision against bacteria.

- **Male dogs** have an incision just above the scrotum. The scrotum is not removed, only the testicles.
- **Male cats** have 1-2 incisions, one on each side of the scrotum, the scrotum is not removed, only the testicles. Male cats may appear as if they still have testicles. This is normal; the swelling should subside gradually through the recovery period. Male cats do not have any sutures.
- **Tattoo** All female cats; female dogs and male dogs receive a small black tattoo either near the incision site or on their umbilicus. This is universally recognized tattoo that identifies your pet has been spayed or neutered.
- **DO NOT GIVE:** Aspirin, Advil, Tylenol or any human medications to your pet for pain relief. *These medications can be deadly.*
- **Complications** Spaying and neutering are both very safe surgeries; however with all surgery, complications can occur. Minimal redness and swelling should resolve within several days, but if they persist longer, please contact us. You should also contact us immediately if you notice any of the following:

Pale Gums
Depression
Vomiting
Diarrhea
Discharge or bleeding from the incision
Difficulty urinating
Labored breathing

The APF will treat at our clinic, at minimal cost, any post-operative complications resulting directly from the surgery, if the above post operative instructions have been followed in full. Your regular veterinarian must address illnesses or injuries that are not a direct result of surgery. We cannot be held responsible for complications resulting from failure to follow post-operative instructions or from contagious diseases for which the animal was not previously properly vaccinated.

In-heat females need to be kept away from males. If your female dog or cat was in heat at the time of surgery, you must keep her away from unneutered males for at least two weeks. While she is now unable to become pregnant, she will still attract intact males for a short period of time. If a male dog attempts to breed a female at this point, it can cause serious, possibly life-threatening damage.

CALL IF YOU HAVE CONCERNS

If you have questions or concerns directly related to the surgery during the recovery period, please call us at 518-374-3944, ext.121 or 125. Please leave a message if you are put through to voicemail and someone will return your call.



If you have an emergency outside of clinic business hours, contact your regular veterinarian or the Capital District Animal Emergency Clinic at 518-785-1094.

Rechecks We can schedule a free recheck for you with our veterinarian if you are concerned about your pet's recovery. You must schedule recheck appointments in advance. We cannot accommodate walk-ins. The APF will provide free rechecks only for surgery related ailments. Any self-inflicted ailments (caused by licking, excessive activity, etc.) are at the owner's expense. Any services rendered by outside veterinarians will not be covered by the APF.